

**Reye's Syndrome:**

Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**

Aspirin may cause a severe allergic reaction, which may include:

- Facial swelling
- Asthma (wheezing)
- Hives
- Shock

**Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- Are age 60 or older
- Have had stomach ulcers or bleeding problems
- Take a blood-thinning (anticoagulant) or steroid drug
- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- Have 3 or more alcoholic drinks every day while using this product
- Take more or for a longer time than directed

**Caffeine Warning:** This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use** if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

**Ask a doctor before use if:**

- Stomach bleeding warning applies to you
- You have a history of stomach problems, such as heartburn

- You have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- You are taking a diuretic
- You have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for:**

- Gout
- Diabetes
- Arthritis

**Stop use and ask a doctor if:**

- An allergic reaction occurs. Seek medical help right away.
- Ringing in ears or loss of hearing occurs
- Any new symptoms occur
- Pain gets worse or lasts more than ten (10) days
- Fever gets worse or lasts more than three (3) days
- Redness or swelling is present
- You experience any of the following signs of stomach bleeding:
  - **Feel faint**
  - **Vomit blood**
  - **Have bloody or black stools**
  - **Have stomach pain that does not get better**

**Each of these could be a sign of a serious condition.**

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin during the last three (3) months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is crucial for adults as well as for children even if you do not notice any signs or symptoms.

**Comments, Questions or Concerns?**

Call toll-free 1-866-255-5197.