

Reye's Syndrome:

Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Aspirin may cause a severe allergic reaction, which may include:

- Hives
- Facial swelling
- Shock
- Asthma (wheezing)

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- Are age 60 or older
- Have had stomach ulcers or bleeding problems
- Take a blood-thinning (anticoagulant) or steroid drug
- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- Have 3 or more alcoholic drinks every day while using this product
- Take more or for a longer time than directed

Caffeine Warning: This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

Ask a doctor before use if:

- Stomach bleeding warning applies to you
- You have a history of stomach problems, such as heartburn

- You have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- You are taking a diuretic
- You have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- Gout
- Diabetes
- Arthritis

Stop use and ask a doctor if:

- An allergic reaction occurs. Seek medical help right away.
- You experience any of the following signs of stomach bleeding:
 - **Feel faint**
 - **Vomit blood**
 - **Have bloody or black stools**
 - **Have stomach pain that does not get better**
- Pain gets worse or lasts more than ten (10) days
- Fever gets worse or lasts more than three (3) days
- Redness or swelling is present
- Any new symptoms occur
- Ringing in ears or loss of hearing occurs

Each of these could be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last three (3) months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs

or symptoms.

Comments, Questions or Concerns?

Call toll-free 1-866-255-5197.